# Gene Therapy for Rare Diseases (muscular dystrophies)



Scott Q. Harper, Ph.D.

Center for Gene Therapy

The Research Institute at Nationwide Children's Hospital,

Department of Pediatrics

The Ohio State University College of Medicine



# Harper Lab Program Goals:

Develop gene therapy approaches for dominantly inherited muscular dystrophies

> RNA interference (RNAi)





### Rare Diseases

- •NIH: diseases affecting less than 200,000 people in U.S.
- •INSERM: 1 in 2,000 Europeans
- •>7,000 rare disorders known; not all are genetic diseases<sup>1,2</sup>
- •Affect 18 25 million Americans (6-8% of population)<sup>1</sup>
- •Maybe affect 300 million people worldwide<sup>3</sup>
- •FY2011: ~9,400 research projects on rare diseases (\$3.5B / 11% of NIH research budget)<sup>4</sup>

<sup>1</sup>NIH Office of Rare Diseases Research





<sup>&</sup>lt;sup>2</sup>Orphanet, INSERM

<sup>&</sup>lt;sup>3</sup>Globalgenes.org/rarelist

<sup>&</sup>lt;sup>4</sup>report.nih.gov/rcdc/categories

### All myopathies are rare disorders





- •Myopathy = muscle disease
- •The Muscular Dystrophy Association (MDA) website lists 91 categories of muscle disease some overlap between categories<sup>1</sup>
- •NINDS groups myopathies into 12 subclasses<sup>2</sup>
- •The Muscular Dystrophies are a group of disorders representing one major subclass of myopathy
- Dominant and recessive forms
- •~50 genes involved in various forms of muscular dystrophy

<sup>1</sup>www.mda.org


<sup>2</sup>www.ninds.nih.gov/disorders/myopathy/myopathy.htm





### Muscle Gene Therapy

- •Emerged from the identification of **Dystrophin** mutations as the cause of X-linked recessive Duchenne Muscular Dystrophy in 1986
  - •DMD is the most common muscular dystrophy (1 in 3,500 newborn males)
  - •~8,000 males in U.S. extrapolates to 1 in 19,000 relative to total U.S. males
  - •First positionally cloned muscular dystrophy disease gene
- •Replacing missing or non-functional *dystrophin* seemed a straightforward DMD treatment (tools were available for



gene replacement)

Dystrophin gene replacement in mdx mouse model of DMD;

Gregorevic, et al Nat Med 2006



replacement



### Historically DMD has been a dominant focus for the muscle gene therapy field

FY2011: 43% of "myopathy/MD" grants on DMD

\$3.6B

NIH expenditures on rare diseases in FY2011



\$75M

- NIH grants on "myopathy/muscular dystrophy" FY2011
- 0.15% of rate disease budget

\$32M

- NIH grants on "DMD" FY2011
  - 43% of \$75M

"Follow the money"



Source: report.nih.gov/categorical\_spending.aspx

FY2011: 85% of muscle gene therapy grants on DMD

\$248M

• NIH grants on "gene therapy" FY2011



\$5.6M

 NIH grants on "gene therapy for myopathy/muscular dystrophy"
2.2% of NIH gene therapy grants



NIH grants related to "DMD gene therapy"





# DMD *gene replacement* focus facilitated muscle gene therapy research in general

- •Advancements in DMD gene therapy are applicable to other muscular dystrophies
  - •Recessive diseases: Successful Phase I α-sarcoglycan gene replacement trial for **LGMD2D** (Mendell, JR et al Ann of Neurol 2008 and 2010)
  - •Dominant diseases: UNEXPLORED UNTIL NOW My lab's focus



Emergence of RNAi provided the tools to treat dominant myopathies with gene therapy

# Dominant Myopathies

- •At least 37 different loci involved in dominant myopathy
- •All rare to extremely rare
- •Silencing of mutant allele may be a common therapeutic strategy
- •Collectively prevalence unclear but may be 1 in 2,400 to 1 in 3,200
  - •Could affect ~130,000 people in U.S.

•My lab is developing therapies for two dominant MDs:

#### **FSHD**

1 in 7,500 – 20,000 16K – 42K Americans Most common dominant MD?





~ 1 in 1,000,000 ~315 Americans Good paradigm for first translation

### LGMD1A

- Age of onset: 18-37 years
- Develop proximal leg and arm weakness in early adulthood and progress to the distal limb
- Myopathic features: variable fiber size, central nuclei, mononuclear infiltration
- Eosinophilic protein aggregates
- Gain-of-function mutation in the Myotilin (MYOT) gene

Myotilin-positive protein aggregates






Olivé M et al. Myotilinopathy: refining the clinical and myopathological phenotype. Brain. 2005;128:2315-26.





### Myotilin

- Expressed primarily in skeletal and cardiac muscle
- Functions as a structural component of the Z-disc





- Not required for normal muscle development or function
- Mouse and human myotilin transcripts have the same expression pattern and are highly conserved





#### **Myotilin T57I Transgenic Mouse Model**





Garvey S M et al. Hum. Mol. Genet. 2006;15:2348-2362






# Hypothesis: Reducing the levels of mutant MYOT will offer a treatment for LGMD1A

Strategy: Develop RNAi gene therapy to suppress mutant MYOT






# Strategy: Deliver MYOT-targeted miRNAs to LGMD1A mice








# AAV.miMYOT mediates efficient MYOT knockdown *in vivo*











# miMYOT Silencing Reduces Protein Aggregates by 70%







# MYOT Silencing Improves Muscle Mass in 3 and 9-month old mice




Overall ~ 10% increase in muscle mass; N=12 mice; p=0.0019





#### MYOT silencing improves whole muscle strength







#### **Conclusions**

- MYOT gene silencing improves muscle abnormalities in mice over-expressing human mutant MYOT
- Moderate expression of miMYOT slows disease progression
- RNAi-based approaches may provide a therapy for LGMD1A and other dominantly inherited myopathies

#### **Ongoing Work**

- Increased dosage of miMYOT may further improve correction (remove GFP)
- Reversal study ongoing
- Preliminary tox studies ongoing
- IP filed





### Hurdles to translation

- •Immunology issues
- •Need good natural history studies
  - "clinically relevant endpoints"
  - •Gene expression/knockdown is not enough must produce functional improvements
  - •Biomarkers must correlate to some functional correction
- •Navigating the patth forward from pre-clinical to IND is confusing and daunting
  - •What are the steps required? Which academic scientists can advise on how to write an IND (~600 pages!)? What's involved in pre-pre-IND, pre-IND meetings with FDA? How do we fund IND-enabling tox, which can run ~\$300K? Where do we produce vector? How do we fund it?
- •Vector Production what do we need for systemic muscle gene therapy? 10<sup>16</sup> particles per person?
  - •1000 liter prep =  $\sim 10^{17}$  = 10 patients: probably max capacity of academic GMP vector core
  - •Estimated \$500k \$1M for one prep
  - •Contract manufacturing needed to provide economy of scale





### Acknowledgements

- Harper Lab Nationwide Children's Hospital
  - Dr. Jian Liu
  - Dr. Lindsay Wallace, Sara Garwick-Coppens
- Collaborators:
  - Jerry Mendell (Nationwide Children's Hospital)
  - Michael Hauser (Duke University)
  - Susan Brooks (University of Michigan)
  - Chady Hakim (University of Missouri)

Other expertise at NCH: Reed Clark, Kevin Flanigan, Chris Schilling, Jerry Mendell

- Funding for LGMD1A pre-clinical work
  - NINDS 1R21NS072260-01
  - Nationwide Children's Hospital





# Gene replacement is not useful for dominantly inherited disorders

